arXiv:1211.3426v4 [math.OA] 29 Jan 2013

Analytic semigroups on vector valued noncommutative L^p-spaces

Cédric Arhancet

Abstract

We give sufficient conditions on an operator space E and on a semigroup of operators on a von Neumann algebra M to obtain a bounded analytic or a R-analytic semigroup $(T_t \otimes Id_E)_{t\geqslant 0}$ on the vector valued noncommutative L^p-space $L^p(M,E)$. Moreover, we give applications to the $H^\infty(\Sigma_\theta)$ functional calculus of the generators of these semigroups, generalizing some earlier work of M. Junge, C. Le Merdy and Q. Xu.

1 Introduction

It is shown in [JMX] that whenever $(T_t)_{t\geqslant 0}$ is a noncommutative diffusion semigroup on a von Neumann algebra M equipped with a faithful normal state such that each T_t satisfies the Rota dilation property, then the negative generator of its L^p-realization ($1<p<\infty$) admits a bounded $H^\infty(\Sigma_\theta)$ functional calculus for some $0<\theta<\frac{\pi}{2}$ where $\Sigma_\theta=\{z\in\mathbb{C}^* : |\arg z|<\theta\}$ is the open sector of angle 2θ around the positive real axis $(0,+\infty)$. Our first principal result is an extension of this theorem to the vector valued case. We provide a different approach using R-analyticity instead of square functions.

In order to describe our result, we need several definitions.

Definition 1.1 *Let (M,ϕ) and (N,ψ) be von Neumann algebras equipped with normal faithful states ϕ and ψ respectively. A linear map $T\colon M\to N$ is called a (ϕ,ψ)-Markov map if*

(1) *T is completely positive*

(2) *T is unital*

(3) $\psi\circ T=\phi$

(4) *$T\circ\sigma_t^\phi=\sigma_t^\psi\circ T$, for all $t\in\mathbb{R}$, where $(\sigma_t^\phi)_{t\in\mathbb{R}}$ and $(\sigma_t^\psi)_{t\in\mathbb{R}}$ denote the automorphism groups of the states ϕ and ψ respectively.*

In particular, when $(M,\phi)=(N,\psi)$, we say that T is a ϕ-Markov map.

A linear map $T\colon M\to N$ satisfying conditions $(1)-(3)$ above is normal. If, moreover, condition (4) is satisfied, then it is known that there exists a unique completely positive, unital map $T^*\colon N\to M$ such that

$$\phi\big(T^*(y)x\big)=\psi\big(yT(x)\big),\qquad x\in M,\ y\in N.$$

The next definition is a variation of the one of [AnD] (see also [Ric] and [HaM]).

The author is supported by the research program ANR 2011 BS01 008 01.
2010 *Mathematics subject classification:* 46L51, 46L07, 47D03.
Key words: noncommutative L^p-spaces, operator spaces, semigroups, functional calculus, $OUMD_p$ property.

Definition 1.2 *A (ϕ,ψ)-Markov map $T\colon M \to N$ is called QWEP-factorizable if there exist a von Neumann algebra P with QWEP equipped with a faithful normal state χ, and $*$-monomorphisms $J_0\colon M \to P$ and $J_1\colon N \to P$ such that J_0 is (ϕ,χ)-Markov and J_1 is (ψ,χ)-Markov, satisfying, moreover, $T = J_0^* \circ J_1$. We say that T is Hyper-factorizable if the same property is true with a hyperfinite von Neumann algebra P.*

Now, we introduce the following definition (compare [HaM, Definition 4.1] and [Arh, Property 4.10])

Definition 1.3 *Let M be a von Neumann algebra equipped with a normal faithful state ϕ. Let $(T_t)_{t\geqslant 0}$ be a w^*-continuous semigroup of ϕ-Markov maps on M. We say that the semigroup is QWEP-dilatable if there exist a von Neumann algebra N with QWEP equipped with a normal faithful state ψ, a w^*-continuous group $(U_t)_{t\geqslant 0}$ of $*$-automorphisms of N, a $*$-monomorphism $J\colon M \to N$ such that each U_t is ϕ-Markov and J is (ϕ,ψ)-Markov satisfying*

$$T_t = \mathbb{E} \circ U_t \circ J, \qquad t \geqslant 0,$$

where $\mathbb{E} = J^*\colon N \to M$ is the canonical faithful normal conditional expectation preserving the states associated with J.

Let $C^*(\mathbb{F}_\infty)$ be the full group C^*-algebra of the free group $\mathbb{F}_\infty$. We say that an operator space E is locally-$C^*(\mathbb{F}_\infty)$ if

$$d_f(E) = \sup_{F\subset E,\text{finite dimensional}} \inf\big\{d_{cb}(F,G) : G \subset C^*(\mathbb{F}_\infty)\big\} < \infty.$$

All natural examples satisfy $d_f(E) = 1$ (see [Pis6, Chapter 21] and [Har] for more information on this class of operator spaces). If M be a von Neumann algebra with *QWEP* equipped with a normal faithful state and if E is locally-$C^*(\mathbb{F}_\infty)$, then the vector valued non commutative L^p-space $L^p(M,E)$ is well-defined and generalize the classical construction for hyperfinite von Neumann algebras (See section 2 for the details).

For any index set I, we denote by $OH(I)$ the associated operator Hilbert space introduced by G. Pisier, see [Pis4] and [Pis6] for more information. Recall that $OUMD_p$ is the operator space analogue of the *Unconditional Martingale Differences* (*UMD*) property of Banach spaces, see section 2 for more information. We also use a similar property $OUMD'_p$ for QWEP von Neumann algebras. The definition is given in section 2. Our main result is the following theorem.

Theorem 1.4 *Let M be a von Neumann algebra with QWEP equipped with a normal faithful state. Let $(T_t)_{t\geqslant 0}$ be a QWEP-dilatable semigroup of QWEP-factorizable maps (resp. Hyper-factorizable maps) on M. Suppose $1 < p,q < \infty$ and $0 < \alpha < 1$. Let E be a locally-$C^*(\mathbb{F}_\infty)$ operator space with $d_f(E) = 1$ such that $E = \big(OH(I),F\big)_\alpha$ with $\frac{1}{p} = \frac{1-\alpha}{2} + \frac{\alpha}{q}$ for some index set I and for some $OUMD'_q$-operator space F (resp. some $OUMD_q$-operator space F if any T_t is Hyper-factorizable). We let $-A_p$ be the generator of the strongly continuous semigroup $(T_t \otimes Id_E)_{t\geqslant 0}$ on $L^p(M,E)$. Then for some $0 < \theta < \frac{\pi}{2}$, the operator A_p has a completely bounded $H^\infty(\Sigma_\theta)$ functional calculus.*

This result can be used with the noncommutative Poisson semigroup or the q-Ornstein-Uhlenbeck semigroup and by example $E = OH(I)$ for any index set I. A version of this result for semigroups of Schur multipliers is also given.

A famous theorem of G. Pisier [Pis2] says that a Banach space X is K-convex (i.e. does not contain uniformly ℓ^1_n's) if and only if the vectorial Rademacher projection $P \otimes Id_X$ is bounded on the Bochner space $L^2(\Omega,X)$ where Ω is a probability space. In this proof, he showed and used the fact that if X is K-convex then any w^*-continuous semigroup $(T_t)_{t\geqslant 0}$ of positive unital selfadjoint Fourier multipliers on a locally compact abelian group G induces a bounded analytic semigroup $(T_t \otimes Id_X)_{t\geqslant 0}$

on the Bochner space $L^p(G,X)$ where $1<p<\infty$. Moreover, he showed a similar result for general w^*-continuous semigroups of positive unital contractions on a measure space if X does not contain, for some $\lambda>1$, any subspace λ-isomorphic to ℓ^1_2. We give noncommutative analogues of these results. They are crucial steps in the proof of our Theorem 1.4.

We say that an operator space E is OK-convex if the vector valued Schatten space $S^2(E)$ is K-convex. This notion was introduced in [JuP]. Using the preservation of the K-convexity under complex interpolation (see [Pis1]), it is easy to see that it is equivalent to the K-convexity of the Banach space $S^p(E)$ for any (all) $1<p<\infty$.

Our second principal result is the following theorem:

Theorem 1.5 *Suppose that G is an amenable discrete group or that G is the free group $\mathbb{F}_n$ with n generators ($1\leqslant n\leqslant\infty$). Let $(T_t)_{t\geqslant 0}$ be a w^*-continuous semigroup of self-adjoint completely positive unital Fourier multipliers on the group von Neumann algebra $VN(G)$ preserving the canonical trace. Let E be a OK-convex operator space. If $G=\mathbb{F}_n$, we suppose that E is locally-$C^*(\mathbb{F}_\infty)$ with $d_f(E)=1$. Consider $1<p<\infty$. Then $(T_t\otimes Id_E)_{t\geqslant 0}$ defines a strongly continuous bounded analytic semigroup on the Banach space $L^p\big(VN(G),E\big)$.*

We will show that this result can be used, by exemple, in the case where E is a Schatten space S^q or a commutative L^q-space with $1<q<\infty$.

The next theorem is a R-analytic version of Theorem 1.5 and it is our last principal result.

Theorem 1.6 *Let M be a von Neumann algebra with $QWEP$ equipped with a normal faithful state. Let $(T_t)_{t\geqslant 0}$ a w^*-continuous semigroup of $QWEP$-factorizable maps (resp. Hyper-factorizable maps) on M. Suppose $1<p,q<\infty$ and $0<\alpha<1$. Let E be a locally-$C^*(\mathbb{F}_\infty)$ operator space with $d_f(E)=1$ such that $E=\big(OH(I),F\big)_\alpha$ with $\frac{1}{p}=\frac{1-\alpha}{2}+\frac{\alpha}{q}$ for some index set I and for some $OUMD'_q$-operator space F (resp. some $OUMD_q$-operator space F if any T_t is Hyper-factorizable). Then $(T_t\otimes Id_E)_{t\geqslant 0}$ defines a strongly continuous R-analytic semigroup on the Banach space $L^p(M,E)$.*

Finally, we also give a version of these two theorems for semigroups of Schur multipliers.

The paper is organized as follows. Section 2 gives a brief presentation of vector valued noncommutative L^p-spaces and we introduce some notions which are relevant to our paper. The next section 3 contains a proof of Theorem 1.5. Section 4 is devoted to prove Theorem 1.6. In section 5, we give applications to functional calculus. In particular, we prove Theorem 1.4. Finally, we present some natural examples to which the results of this paper can be applied.

2 Preliminaries

The readers are referred to [ER], [Pau] and [Pis6] for details on operator spaces and completely bounded maps.

The theory of vector valued noncommutative L^p-spaces was initiated by G. Pisier [Pis5] for the case where the underlying von Neumann algebra is hyperfinite and equipped with a faithful normal semifinite trace. Suppose $1\leqslant p<\infty$. For an operator space E and a hyperfinite von Neumann algebra M equipped with a faithful normal semifinite trace, we define by complex interpolation

$$L^p(M,E)=\big(M\otimes_{\min}E,L^1(M)\widehat{\otimes}E\big)_{\frac{1}{p}}, \tag{2.1}$$

where $\otimes_{\min}$ and $\widehat{\otimes}$ denote the injective and the projective tensor product of operator spaces. In [Jun1] and [Jun2], M. Junge extended this theory to the case where the underlying von Neumann algebra satisfies $QWEP$ using the following characterization of $QWEP$ von Neumann algebras. It is unknown whether every von Neumann algebra has this property. See the survey [Oza] for more information on this notion.

Proposition 2.1 *A von Neumann algebra M is $QWEP$ if and only if there exist a free ultrafilter $\mathcal{U}$ on an index set L and a normal $*$-monomorphism*

$$\pi : M \to B\big(\ell^2_I\big)^{\mathcal{U}}$$

and a normal conditional expectation

$$\mathbb{E}\colon B\big(\ell^2_I\big)^{\mathcal{U}} \to \pi(M)$$

where $B\big(\ell^2_I\big)^{\mathcal{U}}$ denote the ultrapower of $B\big(\ell^2_I\big)$ associated with $\mathcal{U}$.

Note that in general, $\mathbb{E}$ is not faithful.

Now, we introduce the vector valued noncommutative L^p-spaces associated to a von Neumann algebra with $QWEP$ equipped with a faithful normal semifinite weight. Let M, $\alpha = (\pi, \mathbb{E})$ and I be as in Proposition 2.1. Suppose $1 \leqslant p < \infty$. Let E be an operator space. The vector valued Schatten class $S^p_I(E)$ is defined in (2.1) with $M = B\big(\ell^2_I\big)$. Following M. Junge, the vector valued noncommutative L^p-space $L^p(M, \alpha, E)$ is defined as the closure in $S^p_I(E)^{\mathcal{U}}$ of the closed span of

$$\bigcup_{F\subset E,\text{finite dimensional}} S^p_I(F)^{\mathcal{U}}$$

where F runs over all finite dimensional subspaces of E.

The definition of $L^p(M, \alpha, E)$ depends on the choice of α. However, When E is locally-$C^*(\mathbb{F}_\infty)$, the $L^p(M, \alpha, E)$'s are all equivalent allowing the constant $d_f(E)^2$. Thus, we can still say that $L^p(M, \alpha, E)$ does not depend on the choice of α, and thus we will use the more simpler notation $L^p(M, E)$ instead of $L^p(M, \alpha, E)$.

Note the following vector valued extension property of completely positive maps between noncommutative L^p-spaces, see [Pis3] and [Jun2] .

Proposition 2.2

1. *Let M and N be hyperfinite von Neumann algebras equipped with normal faithful semifinite traces and let E be an operator space. Let $T\colon M \to N$ be a trace preserving unital normal positive map. Suppose $1 \leqslant p < \infty$. Then the operator $T \otimes Id_E$ extends to a bounded operator from $L^p(M, E)$ into $L^p(N, E)$ and we have*

$$\|T \otimes Id_E\|_{L^p(M,E)\to L^p(N,E)} \leqslant \|T\|_{L^p(M)\to L^p(N)}.$$

2. *Let M and N be von Neumann algebras with $QWEP$ equipped with normal faithful states ϕ and ψ respectively and let E be a locally-$C^*(\mathbb{F}_\infty)$ operator space. Let $T\colon M \to N$ be a (ϕ, ψ)-Markov map. Suppose $1 \leqslant p < \infty$. Then the operator $T \otimes Id_E$ extends to a bounded operator from $L^p(M, E)$ into $L^p(N, E)$ and we have*

$$\|T \otimes Id_E\|_{L^p(M,E)\to L^p(N,E)} \leqslant d_f(E)\|T\|_{L^p(M)\to L^p(N)}.$$

Suppose that G is a discrete group. We denote by e_G the neutral element of G. We denote by $\lambda_g\colon \ell^2_G \to \ell^2_G$ the unitary operator of left translation by g and $VN(G)$ the von Neumann algebra of G spanned by the λ_g where $g \in G$. It is an finite algebra with normalized trace given by

$$\tau_G(x) = \big\langle \varepsilon_{e_G}, x(\varepsilon_{e_G})\big\rangle_{\ell^2_G}$$

where $(\varepsilon_g)_{g\in G}$ is the canonical basis of ℓ^2_G and $x \in VN(G)$. Recall that if G is amenable then the von Neumann algebra $VN(G)$ is hyperfinite. A Fourier multiplier is a normal linear map $T\colon VN(G) \to VN(G)$ such that there exists a function $\varphi\colon G \to \mathbb{C}$ such that for any $g \in G$ we have $T(\lambda_g) = \varphi_g\lambda_g$. In this case, we denote T by

$$\begin{array}{cccc} M_\varphi\colon & VN(G) & \longrightarrow & VN(G) \\ & \lambda_g & \longmapsto & \varphi_g\lambda_g. \end{array}$$

Let M be a von Neumann algebra equipped with a semifinite normal faithful trace τ. Suppose that $T\colon M \to M$ is a normal contraction. We say that T is selfadjoint if for all $x, y \in M \cap L^1(M)$ we have

$$\tau\big(T(x)y^*\big) = \tau\big(x(Ty)^*\big).$$

In this case, it is not hard to show that the restriction $T|M \cap L^1(M)$ extends to a contraction $T\colon L^1(M) \to L^1(M)$. By complex interpolation, for any $1 \leqslant p \leqslant \infty$, we obtain a contractive map $T\colon L^p(M) \to L^p(M)$. Moreover, the operator $T\colon L^2(M) \to L^2(M)$ is selfadjoint. If $T\colon M \to M$ is actually a normal selfadjoint complete contraction, it is easy to see that the map $T\colon L^p(M) \to L^p(M)$ is completely contractive for any $1 \leqslant p \leqslant \infty$. It is not difficult to show that a contractive Fourier multiplier $M_\varphi\colon VN(G) \to VN(G)$ is selfadjoint if and only if $\varphi\colon G \to \mathbb{C}$ is a real function. Finally, one can prove that a contractive Schur multiplier $M_A\colon B\big(\ell^2_I\big) \to B\big(\ell^2_I\big)$ associated with a matrix A is selfadjoint if and only if all entries of A are real.

Now, we introduce the operator space version of the Banach property UMD. Let E be an operator space and $1 < p < \infty$. We say that E is $OUMD_p$ if there exists a positive constant C such that for any positive integer n, any choice of signs $\varepsilon_k = \pm 1$ and any martingale difference sequence $(dx_k)_{k=1}^n \subset L^p(M, E)$ relative to a filtration $(M_k)_{k \geqslant 1}$ of a hyperfinite von Neumann algebra M we have

$$\left\| \sum_{k=1}^{n} \varepsilon_k dx_k \right\|_{L^p(M,E)} \leqslant C \left\| \sum_{k=1}^{n} dx_k \right\|_{L^p(M,E)}. \tag{2.2}$$

See [Mus] and [Qiu] for more information on this property. We also need a variant of this property for QWEP von Neumann algebras.

Definition 2.3 *Suppose $1 < p < \infty$. Let E be a locally-$C^*(\mathbb{F}_\infty)$ operator space with $d_f(E) = 1$. We say that E is $OUMD'_p$ if there exists a positive constant C such that for any positive integer n, any choice of signs $\varepsilon_k = \pm 1$ and any martingale difference sequence $(dx_k)_{k=1}^n \subset L^p(M, E)$ relative to a filtration $(M_k)_{k \geqslant 1}$ of a QWEP von Neumann algebra M we have the inequality (2.2).*

Let $1 < q, r < \infty$ and $0 < \theta < 1$ be such that $\frac{1}{p} = \frac{1-\alpha}{q} + \frac{\alpha}{r}$. If (E_0, E_1) is a compatible couple of operator spaces, where E_0 is $OUMD_q$ and E_1 is $OUMD_r$, then the complex interpolation operator space $E_\theta = (E_0, E_1)_\alpha$ is $OUMD_p$. For any index set I and any $1 < p < \infty$, the operator Hilbert space $OH(I)$ is $OUMD_p$. If E is $OUMD_p$ then the Banach space $S^p(E)$ is UMD (hence K-convex). It is easy to see that the same properties are valid for $OUMD'_p$ operator spaces (with the same proofs).

3 Analyticity

Let X be a Banach space. A strongly continuous semigroup $(T_t)_{t \geqslant 0}$ is called bounded analytic if there exist $0 < \theta < \frac{\pi}{2}$ and a bounded holomorphic extension

$$\begin{array}{ccc} \Sigma_\theta & \longrightarrow & B(X) \\ z & \longmapsto & T_z. \end{array}$$

See [KW] and [Haa] for more information on this notion.

We need the following theorem which is a corollary [Pis2, Lemma 4] of a result of Beurling [Beu] (see also [Fac]).

Theorem 3.1 *Let X be a Banach space. Let $(T_t)_{t \geqslant 0}$ a strongly continuous semigroup of contractions on X. Suppose that there exists some integer $n \geqslant 1$ such that for any $t > 0$*

$$\big\| (Id_X - T_t)^n \big\|_{X \to X} < 2^n.$$

Then the semigroup $(T_t)_{t \geqslant 0}$ is bounded analytic.

Moreover, we will use the following lemma [Pis2, Lemma 1.5].

Lemma 3.2 *Let $n \geqslant 1$ be an integer. Suppose that X is a real Banach space, such that, for some $\lambda > 1$, X does not contain any subspace λ-isomorphic to ℓ^1_{n+1}. Then there exists a real number $0 < \rho < 2$ such that if $P_1, \ldots, P_n$ is any finite collection of mutually commuting norm one projections on X, then*

$$\Bigg\| \prod_{1 \leqslant k \leqslant n} (Id_X - P_k) \Bigg\|_{X \to X} \leqslant \rho^n.$$

Note that it is known that a complex Banach space contains ℓ^1_n's uniformly if and only if the underlying real Banach space has the same property.

Finally, we recall the next result ([DH, Lemma 1].

Lemma 3.3 *Let M and N be two von Neumann algebras equipped with faithful normal finite traces, and let $(x_i)_{i\in I} \in M$ and $(y_i)_{i \in I} \in N$ be families of elements of M and N respectively that have the same $*$-distribution. Then the von Neumann algebras generated respectively by the x_i's and the y_i's are isomorphic, with a normal $*$-isomorphism sending x_i on y_i and preserving the trace.*

We will use the next result which is a variant of well-known Fell's absorption principle (see [Pis6, Proposition 8.1]). This proposition is a substitute for the trick of G. Pisier [Pis2, Lemma 1.6].

Proposition 3.4 *Let G be a discrete group and $1 \leqslant p < \infty$. Let E be an operator space. If G is non-amenable, we suppose that E is locally-$C^*(\mathbb{F}_\infty)$ with $d_f(E) = 1$. For any function $a \colon G \to E$ finitely supported on G, we have*

$$\Bigg\| \sum_{g \in G} \lambda_g \otimes \lambda_g \otimes \cdots \otimes \lambda_g \otimes a(g) \Bigg\|_{L^p(\overline{\otimes}_{i=1}^n VN(G), E)} = \Bigg\| \sum_{g \in G} \lambda_g \otimes a(g) \Bigg\|_{L^p(VN(G),E)}. \tag{3.1}$$

Moreover, for any completely positive unital Fourier multiplier $M_\varphi \colon VN(G) \to VN(G)$ preserving the canonical trace and any positive integer n we have

$$\big\| M_\varphi^n \otimes Id_E \big\|_{L^p(VN(G),E) \to L^p(VN(G),E)} \leqslant \big\| M_\varphi^{\otimes n} \otimes Id_E \big\|_{L^p(\overline{\otimes}_{i=1}^n VN(G),E) \to L^p(\overline{\otimes}_{i=1}^n VN(G),E)}. \tag{3.2}$$

Proof : Suppose that m is an integer and that $g_{r_1}, \ldots, g_{r_m} \in G$. Let $\eta_1, \ldots, \eta_m \in \{*, 1\}$ and $\varepsilon_1, \ldots, \varepsilon_m \in \{-1, 1\}$ the associated sign (i.e $\varepsilon_i = -1$ if and only if $\eta_i = *$). For any integer n, we have

$$\begin{aligned} \tau_G^{\otimes n}\Big(\big(\lambda_{g_{r_1}} \otimes \cdots \otimes \lambda_{g_{r_1}}\big)^{\eta_1} \cdots \big(\lambda_{g_{r_m}} \otimes \cdots \otimes \lambda_{g_{r_m}}\big)^{\eta_m} \Big) &= \tau_G\big(\lambda_{g_{r_1}^{\varepsilon_1} \cdots g_{r_m}^{\varepsilon_m}}\big)^n \\ &= \tau_G\big(\lambda_{g_{r_1}^{\varepsilon_1} \cdots g_{r_m}^{\varepsilon_m}}\big) \\ &= \tau_G\Big((\lambda_{g_{r_1}}^{\eta_1} \cdot \cdots \cdot \lambda_{g_{r_m}}^{\eta_m} \Big). \end{aligned}$$

We infer that the families $\big(\lambda_g \otimes \cdots \otimes \lambda_g\big)_{g \in G}$ and $\big(\lambda_g\big)_{g\in G}$ have the same $*$-distribution with respect to the von Neumann algebras $\overline{\otimes}_{i=1}^n VN(G)$ equipped with $\tau_G^{\otimes n}$ and $VN(G)$ equipped with τ_G. We conclude by using Lemma 3.3 and Proposition 2.2.

Now, we prove the second part of the proposition. Using (3.1) twice, for any positive integer n and any function $a\colon G \to E$ finitely supported on G, we have

$$
\begin{aligned}
&\bigg\|(M_\varphi^n \otimes Id_E)\bigg(\sum_{g\in G} \lambda_g \otimes a(g)\bigg)\bigg\|_{L^p(VN(G),E)} = \bigg\|\sum_{g\in G} \varphi_g^n \lambda_g \otimes a(g)\bigg\|_{L^p(VN(G),E)}\\
&\quad = \bigg\|\sum_{g\in G} \varphi_g^n \lambda_g \otimes \cdots \otimes \lambda_g \otimes a(g)\bigg\|_{L^p(\overline{\otimes}_{i=1}^n VN(G),E)}\\
&\quad = \bigg\|(M_\varphi \otimes \cdots \otimes M_\varphi \otimes Id_E)\bigg(\sum_{t\in G} \lambda_g \otimes \cdots \otimes \lambda_g \otimes a(g)\bigg)\bigg\|_{L^p(\overline{\otimes}_{i=1}^n VN(G),E)}\\
&\quad \leqslant \big\|M_\varphi \otimes \cdots \otimes M_\varphi \otimes Id_E\big\|_{L^p(\overline{\otimes}_{i=1}^n VN(G),E) \to L^p(\overline{\otimes}_{i=1}^n VN(G),E)} \bigg\|\sum_{g\in G} \lambda_g \otimes \cdots \otimes \lambda_g \otimes a(g)\bigg\|_{L^p(\overline{\otimes}_{i=1}^n VN(G),E)}\\
&\quad = \big\|M_\varphi \otimes \cdots \otimes M_\varphi \otimes Id_E\big\|_{L^p(\overline{\otimes}_{i=1}^n VN(G),E) \to L^p(\overline{\otimes}_{i=1}^n VN(G),E)} \bigg\|\sum_{g\in G} \lambda_g \otimes a(g)\bigg\|_{L^p(VN(G),E)}.
\end{aligned}
$$

We deduce (3.2). ■

Proposition 3.5 *Let M be a von Neumann with QWEP equipped with a normal semifinite faithful weight. Let E be a locally-$C^*(\mathbb{F}_\infty)$ operator space which is OK-convex. Suppose $1 < p < \infty$. Then the Banach space $L^p(M,E)$ is K-convex.*

Proof : By definition, the Banach space $L^p(M,E)$ is a closed subspace of a ultrapower $S_I^p(E)^{\mathcal{U}}$ of a vector valued Schatten space $S_I^p(E)$ for some index set I. The Banach space $S^p(E)$ is K-convex. Hence the Banach space $S_I^p(E)$ is also K-convex. Recall that K-convexity is a super-property [DJT, page 261], i.e. passes from a Banach space to all closed subspaces of its ultrapowers. We conclude that the Banach space $L^p(M,E)$ is K-convex. ■

Now, we are ready to give the proof of Theorem 1.5.

Proof of Theorem 1.5 : By [Ric], for any $t \geqslant 0$, the operator $T_t = (T_{\frac{t}{2}})^2$ admits a Rota dilation (see [HaM, Definition 5.1] and [JMX, page 124] for a precise definition)

$$T_t^k = Q\mathbb{E}_k\pi, \qquad k \geqslant 1$$

where $Q\colon M \to VN(G)$ and $\pi\colon VN(G) \to M$ and where we use a crossed product

$$M = \Gamma_{-1}\big(\ell^{2,T} \otimes_2 \ell^2\big) \rtimes_\alpha G$$

equipped with its canonical trace. We infer that

$$Id_{L^p(VN(G))} - T_t = Q(Id_{L^p(M)} - \mathbb{E}_1)\pi.$$

Let n be a positive integer. We deduce that

$$(Id_{L^p(VN(G))} - T_t)^{\otimes n} = Q(Id_{L^p(M)} - \mathbb{E}_1)^{\otimes n}\pi.$$

For any integer $1 \leqslant j \leqslant n$, we let

$$\Pi_j = Id_{L^p(M)} \otimes \cdots \otimes Id_{L^p(M)} \otimes \mathbb{E}_1 \otimes Id_{L^p(M)} \otimes \cdots \otimes Id_{L^p(M)}.$$

Recall that the fermion algebra $\Gamma_{-1}\big(\ell^{2,T} \otimes_2 \ell^2\big)$ is $*$-isomorphic to the hyperfinite factor of type II_1. Moreover, if G is amenable then by [Con, Proposition 6.8], M is hyperfinite. If $G = \mathbb{F}_n$, by [Arh, Proposition 4.8], the von Neumann algebra M has $QWEP$. By Proposition 2.2, we deduce that the $\Pi_j \otimes Id_E$'s are well-defined and form a family of mutually commuting contractive projections on $L^p(M,E)$. Moreover, we have

$$\big((Id_{L^p(M)} - \mathbb{E}_1)\big)^{\otimes n} \otimes Id_E = \prod_{1 \leqslant j \leqslant n} \Big(Id_{L^p(M,E)} - (\Pi_j \otimes Id_E)\Big).$$

Using (3.2), Proposition 2.2 and Lemma 3.2, we obtain that

$$\begin{aligned}
&\Big\|\big(Id_{L^p(VN(G))} - T_t\big)^n \otimes Id_E\Big\|_{L^p(VN(G),E) \to L^p(VN(G),E)} \\
&\leqslant \Big\|(Id_{L^p(VN(G))} - T_t)^{\otimes n} \otimes Id_E\Big\|_{L^p(\overline{\otimes}_{i=1}^n VN(G),E) \to L^p(\overline{\otimes}_{i=1}^n VN(G),E)} \\
&= \Big\|Q(Id_{L^p(M)} - \mathbb{E}_1)^{\otimes n}\pi \otimes Id_E\Big\|_{L^p(\overline{\otimes}_{i=1}^n VN(G),E) \to L^p(\overline{\otimes}_{i=1}^n VN(G),E)} \\
&\leqslant \Big\|(Id_{L^p(M)} - \mathbb{E}_1)^{\otimes n} \otimes Id_E\Big\|_{L^p(M,E) \to L^p(M,E)} \\
&= \Bigg\|\prod_{1 \leqslant j \leqslant n} \Big(Id_{L^p(M,E)} - (\Pi_j \otimes Id_E)\Big)\Bigg\|_{L^p(M,E) \to L^p(M,E)} \\
&\leqslant \rho^n.
\end{aligned}$$

We conclude by applying Theorem 3.1. ■

Now, we pass to general semigroups on $QWEP$ von Neumann algebras. Note that the class of operators spaces considered in the following theorem is smaller than the class of OK-convex operator spaces.

Theorem 3.6 *Let M a von Neumann algebra with $QWEP$ equipped with a normal faithful state. Let $(T_t)_{t \geqslant 0}$ be a w^*-continuous semigroup of $QWEP$-factorizable maps on M. Suppose $1 < p < \infty$. Let E be a locally-$C^*(\mathbb{F}_\infty)$ operator space with $d_f(E) = 1$ such that $S^p(E)$ does not contain, for some $\lambda > 1$, any subspace λ-isomorphic to ℓ_1^2. Then $(T_t \otimes Id_E)_{t \geqslant 0}$ defines a strongly continuous bounded analytic semigroup on the Banach space $L^p(M,E)$.*

Proof : Using the fact that each T_t is $QWEP$-factorizable and [HaM, Theorem 5.3], it is easy to see that each T_t admits a Rota dilation

$$T_t^k = Q\mathbb{E}_k\pi, \qquad k \geqslant 1$$

where $Q\colon M \to N$ and $\pi\colon M \to N$ where N is a $QWEP$ von Neumann algebra. The rest of the proof is similar to the one of Theorem 1.5. The only difference is that we does not use Proposition 3.4. We use Lemma 3.2 with $n = 1$. ■

For instance, if $S^p(E)$ (where $1 < p < \infty$) is uniformly convex then for some $\lambda > 1$, $S^p(E)$ does not contain any subspace λ-isomorphic to ℓ_2^1.

Suppose $1 < p, q < \infty$. Note that if an operator space E is $OUMD_q$ then the Banach space $S^q(E)$ is UMD, hence uniformly convex. Now, we can write $\frac{1}{p} = \frac{1-\alpha}{q} + \frac{\alpha}{r}$ for some $0 < \alpha < 1$ and some $1 < r < \infty$. Then we have $S^p(E) = \big(S^q(E), S^r(E)\big)_\alpha$. Now recall that, by [CwR], if (X_0, X_1) is a compatible couple of Banach spaces, one of which is uniformly convex, then for any $0 < \alpha < 1$ the complex interpolation $(X_0, X_1)_\alpha$ is also uniformly convex. We deduce that the Banach space $S^p(E)$ is uniformly convex. Hence, we can use in Theorem 3.6 (and also in Theorem 1.5) any operator space E which is $OUMD_q$ for *some* $1 < q < \infty$ and locally-$C^*(\mathbb{F}_\infty)$ with $d_f(E) = 1$. This large class contains in particular Schatten spaces S^q and commutative L^q-spaces for any $1 < q < \infty$,

Finally, we deal with semigroups of Schur multipliers. Note that the construction of the Rota Dilation for Schur multipliers on finite dimensional $B(\ell_I^2)$ of [Ric] is actually true for any index set

I. Moreover, the von Neumann algebra $\Gamma^{e}_{-1}(\ell^{2,T})$ of [Ric] is hyperfinite. Hence, the von Neumann algebra

$$M = B\big(\ell^2_I\big)\overline{\otimes}\Bigg(\overline{\bigotimes_{n\in\mathbb{N}}}\big(\Gamma^{e}_{-1}(\ell^{2,T}),\tau\big)\Bigg).$$

of the Rota Dilation of [Ric] is also hyperfinite. Using the above ideas, one can prove the following theorem.

Theorem 3.7 *Let $(T_t)_{t\geqslant 0}$ be a w^*-continuous semigroup of completely positive unital selfadjoint Schur multipliers on $B\big(\ell^2_I\big)$. Suppose $1<p<\infty$. Let E be an operator space such that $S^p(E)$ does not contain, for some $\lambda>1$, any subspace λ-isomorphic to ℓ^2_1. Then $(T_t\otimes Id_E)_{t\geqslant 0}$ defines a strongly continuous bounded analytic semigroup on the Banach space $S^p_I(E)$.*

4 R-analyticity

Let $(\varepsilon_k)_{k\geqslant 1}$ be a sequence of independent Rademacher variables on some probability space $(\Omega,\mathcal{F},\mathbb{P})$. We let $\mathrm{Rad}(X)\subset L^2(\Omega,X)$ be the closure of $\mathrm{Span}\big\{\varepsilon_k\otimes x\,:\,k\geqslant 1,\ x\in X\big\}$ in the Bochner space $L^2(\Omega,X)$. Thus for any finite family $x_1,\ldots,x_n$ in X, we have

$$\Bigg\|\sum_{k=1}^{n}\varepsilon_k\otimes x_k\Bigg\|_{\mathrm{Rad}(X)}=\Bigg(\int_\Omega\Bigg\|\sum_{k=1}^{n}\varepsilon_k(\omega)\,x_k\Bigg\|_X^2 d\omega\Bigg)^{\frac{1}{2}}.$$

We say that a set $F\subset B(X)$ is R-bounded provided that there is a constant $C\geqslant 0$ such that for any finite families $T_1,\ldots,T_n$ in F and $x_1,\ldots,x_n$ in X, we have

$$\Bigg\|\sum_{k=1}^{n}\varepsilon_k\otimes T_k(x_k)\Bigg\|_{\mathrm{Rad}(X)}\leqslant C\Bigg\|\sum_{k=1}^{n}\varepsilon_k\otimes x_k\Bigg\|_{\mathrm{Rad}(X)}.$$

In this case, we let $R(F)$ denote the smallest possible C, which is called the R-bound of F. Note that any singleton $\{T\}$ is automatically R-bounded with $R\big(\{T\}\big)=\|T\|_{X\to X}$. If a set F is R-bounded then the strong closure of F is R-bounded (with the same R-bound). Recall that if H is a Hilbert space, a subset F of $B(H)$ is bounded if and only if F is R-bounded.

R-boundedness was introduced in [BeG] and then developed in the fundamental paper [ClP]. We refer to the latter paper and to [KW, Section 2] for a detailed presentation.

The next result is a noncommutative version of the classical result of Bourgain [Bou] which itself is a vector valued generalization of a result of Stein [Ste].

Proposition 4.1 *Let M be a von Neumann algebra with QWEP equipped with a normal faithful semifinite weight. Suppose $1<p<\infty$. Let E be a $OUMD'_p$-operator space (resp. $OUMD_p$-operator space if M is hyperfinite). Consider a increasing (or decreasing) sequence $\big(\mathbb{E}(\cdot|M_i)\big)_{i\in\mathbb{N}}$ of (canonical) conditional expectations on some von Neumann subalgebras M_i of M. Then the set of conditional expectation operators $\big\{\mathbb{E}(\cdot|M_i)\otimes Id_E\,:\,i\in\mathbb{N}\big\}$ is R-bounded on $L^p(M,E)$.*

Proof : We only prove the decreasing case $M_1\supset M_2\supset\cdots$ and the hyperfinite case. The proofs of other statements are similar. First suppose that the weight is a finite trace. Let n be a positive integer and fix some positive integers $i_1>i_2>\cdots>i_n$. We define the σ-algebra $\mathcal{F}_k=\sigma(\varepsilon_1,\ldots,\varepsilon_k)$, $k=1,\ldots,n$ and $\mathcal{F}_0=\emptyset$. We define the family $(N_m)_{m=1}^{2n}$ of von Neumann subalgebras of $L^\infty(\Omega)\otimes M$ by

$$N_{2k-1}=L^\infty(\Omega,\mathcal{F}_{k-1},\mathbb{P})\otimes M_{i_k},\qquad k=1,\ldots,n$$

$$N_{2k}=L^\infty(\Omega,\mathcal{F}_k,\mathbb{P})\otimes M_{i_k},\qquad k=1,\ldots,n.$$

These subalgebras form an increasing sequence $N_1 \subset N_2 \subset \cdots \subset N_{2n}$. For an element $y \in L^p\big(L^\infty(\Omega) \otimes M, E\big)$, we define the martingale $(y_m)_{1 \leqslant m \leqslant 2n}$ by

$$y_m = \big(\mathbb{E}(\cdot|N_m) \otimes Id_E\big)(y), \qquad m = 1, \ldots, 2n.$$

Let $(d_m)_{1 \leqslant m \leqslant 2n}$ be the associated martingale difference sequence. Then, by the $OUMD_p$-property of E, we have

$$\begin{aligned}
\Big\| \sum_{k=1}^n d_{2k} \Big\|_{L^p(L^\infty(\Omega) \otimes M, E)} &= \frac{1}{2} \Big\| \sum_{m=1}^{2n} d_m + \sum_{m=1}^{2n} (-1)^m d_m \Big\|_{L^p(L^\infty(\Omega) \otimes M, E)} \\
&\leqslant \frac{1}{2} \bigg(\Big\| \sum_{m=1}^{2n} d_m \Big\|_{L^p(L^\infty(\Omega) \otimes M, E)} + \Big\| \sum_{m=1}^{2n} (-1)^m d_m \Big\|_{L^p(L^\infty(\Omega) \otimes M, E)} \bigg) \\
&\lesssim \Big\| \sum_{m=1}^{2n} d_m \Big\|_{L^p(L^\infty(\Omega) \otimes M, E)}
\end{aligned}$$

(here and in the sequel of the proof, we use $\lesssim$ to indicate an inequality up to a constant). Now fix $x_1, \ldots, x_n \in L^p(M, E)$ and put $y = \sum_{l=1}^n \varepsilon_l \otimes x_l$. For this choice of y and any integer $1 \leqslant k \leqslant n$, we have

$$\begin{aligned}
y_{2k-1} &= \big(\mathbb{E}(\cdot|N_{2k-1}) \otimes Id_E\big)(y) \\
&= \sum_{l=1}^n \big(\mathbb{E}\big(\,\cdot\,|L^\infty(\Omega, \mathcal{F}_{k-1}, \mathbb{P}) \otimes M_{i_k}\big) \otimes Id_E\big)(\varepsilon_l \otimes x_l) \\
&= \sum_{l=1}^n \mathbb{E}\big(\varepsilon_l | L^\infty(\Omega, \mathcal{F}_{k-1}, \mathbb{P})\big) \otimes \big(\mathbb{E}(\cdot|M_{i_k}) \otimes Id_E\big)(x_l) \\
&= \sum_{l=1}^{k-1} \varepsilon_l \otimes \big(\mathbb{E}(\cdot|M_{i_k}) \otimes Id_E\big)(x_l)
\end{aligned}$$

and similarly for any integer $1 \leqslant k \leqslant n$ we have

$$y_{2k} = \sum_{l=1}^{k} \varepsilon_l \otimes \big(\mathbb{E}(\cdot|M_{i_k}) \otimes Id_E\big)(x_l).$$

Therefore, for any $1 \leqslant k \leqslant n$, we have

$$\begin{aligned}
d_{2k} &= y_{2k} - y_{2k-1} \\
&= \sum_{l=1}^{k} \varepsilon_l \otimes \big(\mathbb{E}(\cdot|M_{i_k}) \otimes Id_E\big)(x_l) - \sum_{k=1}^{k-1} \varepsilon_l \otimes \big(\mathbb{E}(\cdot|M_{i_k}) \otimes Id_E\big)(x_l) \\
&= \varepsilon_k \otimes \big(\mathbb{E}(\cdot|M_{i_k}) \otimes Id_E\big)(x_k)
\end{aligned}$$

and similarly $d_{2k-1} = 0$ for any integer $1 \leqslant k \leqslant n$. Finally, we obtain that

$$\begin{aligned}
\Big\| \sum_{k=1}^{n} \varepsilon_k \otimes \big(\mathbb{E}(\cdot|M_{i_k}) \otimes Id_E\big)(x_k) \Big\|_{L^p(L^\infty(\Omega)\otimes M,E)} &= \Big\| \sum_{k=1}^{n} d_{2k} \Big\|_{L^p(L^\infty(\Omega)\otimes M,E)} \\
&\lesssim \Big\| \sum_{m=1}^{2n} d_m \Big\|_{L^p(L^\infty(\Omega)\otimes M,E)} \\
&= \big\| \big(\mathbb{E}(\cdot|N_{2n}) \otimes Id_E\big)(y) \big\|_{L^p(L^\infty(\Omega)\otimes M,E)} \\
&\lesssim \Big\| \sum_{k=1}^{n} \varepsilon_k \otimes x_k \Big\|_{L^p(L^\infty(\Omega)\otimes M,E)}.
\end{aligned}$$

By using Khintchine-Kahane inequalities (e.g. [DJT, page 211]), we conclude that

$$\Big\| \sum_{k=1}^{n} \varepsilon_k \otimes \big(\mathbb{E}(\cdot|M_{i_k}) \otimes Id_E\big)(x_k) \Big\|_{\mathrm{Rad}(L^p(M,E))} \lesssim \Big\| \sum_{k=1}^{n} \varepsilon_k \otimes x_k \Big\|_{\mathrm{Rad}(L^p(M,E))}.$$

We deduce the general case of von Neumann algebras equipped with a faithful normal semifinite weight by a straightforward application of the well-known reduction method of Haagerup [HJX]. ■

Let X be a Banach space. A strongly continuous semigroup $(T_t)_{t\geqslant 0}$ is called R-analytic if there exist $0 < \theta < \frac{\pi}{2}$ and an holomorphic extension

$$\begin{array}{ccc} \Sigma_\theta & \longrightarrow & B(X) \\ z & \longmapsto & T_z \end{array}$$

with $R\big(\{T(z) : z \in \Sigma_\theta\}\big) < \infty$. See [KW] for more information and for applications to maximal regularity.

The following result is a particular case of [Fac, Theorem 6.1].

Theorem 4.2 *Let $(T_{1,t})_{t\geqslant 0}$ and $(T_{2,t})_{t\geqslant 0}$ be two consistent semigroups given on an interpolation couple (X_1, X_2) of K-convex Banach spaces. Suppose $0 < \theta < 1$. Assume that $(T_{1,t})_{t\geq 0}$ is strongly continuous and R-analytic and that $R\big(\{T_{2,t} : 0 < t < 1\}\big) < \infty$. Then there exists a unique strongly continuous R-analytic semigroup $(T_t)_{t\geqslant 0}$ on $(X_1, X_2)_\theta$ which is consistent with $(T_{1,t})_{t\geqslant 0}$ and $(T_{2,t})_{t\geqslant 0}$.*

Now, we prove Theorem 1.6.

Proof of Theorem 1.6 : We only prove the Hyper-factorizable case. The QWEP-factorizable case is similar. We can identify $OH(I)$ with $\ell^2(I)$ completely isometrically. For any $i \in I$, we set $E_i = L^2(M)$. Fubini's theorem gives us an isometric isomorphism

$$L^2\big(M, OH(I)\big) = \ell^2\big(\{E_i, i \in I\}\big).$$

Hence the Banach space $L^2\big(M, OH(I)\big)$ is isometric to a Hilbert space. On a Hilbert space, recall that any bounded set is R-bounded. By Theorem 3.6, we deduce that $(T_t \otimes Id_{OH(I)})_{t\geqslant 0}$ defines a R-analytic semigroup on $L^p\big(M, OH(I)\big)$.

Let $t_1, \ldots, t_n$ be rational numbers. We take a common denominator: we can write $t_i = \frac{s_i}{d}$ for some integers $d, s_1, \ldots, s_n$. The operator $T_{\frac{1}{d}}$ admits a Rota dilation:

$$T_{\frac{1}{d}}^k = Q\mathbb{E}_k\pi, \qquad k \geqslant 1.$$

Let $x_1, \ldots, x_n \in F$. Using Proposition 4.1 and the fact that any singleton is R-bounded, we obtain

$$\begin{aligned}
\Bigg\| \sum_{i=1}^{n} \varepsilon_i \otimes (T_{t_i} \otimes Id_F)(x_i) \Bigg\|_{\mathrm{Rad}(L^p(M,F)} &= \Bigg\| \sum_{i=1}^{n} \varepsilon_i \otimes (Q\mathbb{E}_{s_i}\pi \otimes Id_F)(x_i) \Bigg\|_{\mathrm{Rad}(L^p(M,F)} \\
&\leqslant \Bigg\| \sum_{i=1}^{n} \varepsilon_i \otimes (\mathbb{E}_{s_i}\pi \otimes Id_F)(x_i) \Bigg\|_{\mathrm{Rad}(L^p(N,F))} \\
&\leqslant R\big\{\mathbb{E}_k \otimes Id_F \ :\ k \geqslant 1\big\} \Bigg\| \sum_{i=1}^{n} \varepsilon_i \otimes (\pi \otimes Id_F)x_i \Bigg\|_{\mathrm{Rad}(L^p(M,F))} \\
&\leqslant R\big\{\mathbb{E}_k \otimes Id_F \ :\ k \geqslant 1\big\} \Bigg\| \sum_{i=1}^{n} \varepsilon_i \otimes x_i \Bigg\|_{\mathrm{Rad}(L^p(M,F))}.
\end{aligned}$$

By using the strong continuity of the semigroup, We conclude that $\{T_t \otimes Id_F \ :\ t \geqslant 0\}$ is a R-bounded subset of $B\big(L^p(M,F)\big)$.

Now, we have

$$L^p(M,E) = \Big(L^2\big(M, OH(I)\big), L^q(M,F)\Big)_{\alpha}.$$

Furthermore, the operator space F is $OUMD_p$. We deduce that the Banach space $S^p(F)$ is UMD, hence K-convex. Thus the operator space F is OK-convex. By Proposition 3.5, we deduce that $L^q(M,F)$ is K-convex and moreover the Banach space $L^2\big(M, OH(I)\big)$ is obviously K-convex. We conclude by using Theorem 4.2. ■

By example, we can take $E = OH(I)$ in Theorem 1.6.

5 Applications to functional calculus

We start with a little background on sectoriality and H^∞ functional calculus. We refer to [Haa], [KW], [JMX] and [LM] for details and complements. A closed, densely defined linear operator $A\colon D(A) \subset X \to X$ is called sectorial of type ω if its spectrum $\sigma(A)$ is included in the closed sector $\overline{\Sigma_\omega}$, and for any angle $\omega < \theta < \pi$, there is a positive constant K_θ such that

$$\big\|(\lambda - A)^{-1}\big\|_{X \to X} \leqslant \frac{K_\theta}{|\lambda|}, \qquad \lambda \in \mathbb{C} - \overline{\Sigma_\theta}.$$

We recall that sectorial operators of type $< \frac{\pi}{2}$ coincide with negative generators of bounded analytic semigroups.

For any $0 < \theta < \pi$, let $H^\infty(\Sigma_\theta)$ be the algebra of all bounded analytic functions $f\colon \Sigma_\theta \to \mathbb{C}$, equipped with the supremum norm $\|f\|_{H^\infty(\Sigma_\theta)} = \sup\big\{|f(z)| \ :\ z \in \Sigma_\theta\big\}$. Let $H_0^\infty(\Sigma_\theta) \subset H^\infty(\Sigma_\theta)$ be the subalgebra of bounded analytic functions $f\colon \Sigma_\theta \to \mathbb{C}$ for which there exist $s, c > 0$ such that $|f(z)| \leqslant c|z|^s\big(1 + |z|\big)^{-2s}$ for any $z \in \Sigma_\theta$. Given a sectorial operator A of type $0 < \omega < \pi$, a bigger angle $\omega < \theta < \pi$, and a function $f \in H_0^\infty(\Sigma_\theta)$, one may define a bounded operator $f(A)$ by means of a Cauchy integral (see e.g. [Haa, Section 2.3] or [KW, Section 9]); the resulting mapping $H_0^\infty(\Sigma_\theta) \to B(X)$ taking f to $f(A)$ is an algebra homomorphism. By definition, A has a bounded $H^\infty(\Sigma_\theta)$ functional calculus provided that this homomorphism is bounded, that is, there exists a positive constant C such that $\big\|f(A)\big\|_{X \to X} \leqslant C\|f\|_{H^\infty(\Sigma_\theta)}$ for any $f \in H_0^\infty(\Sigma_\theta)$. In the case when A has a dense range, the latter boundedness condition allows a natural extension of $f \mapsto f(A)$ to the full algebra $H^\infty(\Sigma_\theta)$.

Suppose $1 < p < \infty$. In the sequel, we say that an operator A on a vector valued noncommutative L^p-space $L^p(M,E)$ admits a completely bounded $H^\infty(\Sigma_\theta)$ functional calculus if $Id_{S^p} \otimes A$ admits a bounded $H^\infty(\Sigma_\theta)$ functional calculus on the Banach space $S^p\big(L^p(M,E)\big)$.

We will use the following theorem [KW, Corollary 10.9]:

Theorem 5.1 *Let* $(T_t)_{t \geqslant 0}$ *be a strongly continuous semigroup with generator* $-A$ *on a Banach space. If the semigroup has a dilation to a bounded strongly continuous group on a* UMD *Banach space, then* A *has a bounded* $H^\infty(\Sigma_\theta)$ *functional calculus for some* $0 < \theta < \pi$.

We also need the particular case of [KaW, Proposition 5.1] which says that the presence of R-analyticity allows to reduce the angle of a bounded $H^\infty(\Sigma_\theta)$ functional calculus below $\frac{\pi}{2}$.

Proposition 5.2 *Let* $(T_t)_{t \geqslant 0}$ *be a strongly continuous semigroup with generator* $-A$ *on a Banach space. Suppose that* A *has a bounded* $H^\infty(\Sigma_{\theta_0})$ *functional calculus for some* $0 < \theta_0 < \pi$. *If the semigroup* $(T_t)_{t \geqslant 0}$ *is* R*-analytic then* A *has indeed a bounded* $H^\infty(\Sigma_\theta)$ *functional calculus for some* $0 < \theta < \frac{\pi}{2}$.

Now, we give the proof of our main result.

Proof of Theorem 1.4 : By definition, there exists a dilation of the semigroup $(T_t)_{t \geqslant 0}$. This dilation induces a dilation of the semigroup $(T_t \otimes Id_E)_{t \geqslant 0}$ on $L^p(M, E)$ to a strongly continuous group of isometries on $L^p(N, E)$.

Since the operator Hilbert space $OH(I)$ has $OUMD_2$ and the operator space F have $OUMD_q$, we see by interpolation that the operator space E has $OUMD_p$. We infer that the Banach space $S^p(E)$ is UMD. Hence the Banach space $S^p_I(E)$ is also UMD for any index set I. Now, the Banach space $L^p(N, E)$ is a closed subspace of an ultrapower $S^p_I(E)^{\mathcal{U}}$ of a vector valued Schatten space $S^p_I(E)$ for some index set I. We deduce that the Banach space $L^p(N, E)$ is UMD.

By Theorem 5.1, we deduce that A_p has a bounded $H^\infty(\Sigma_\theta)$ functional calculus for some $0 < \theta < \pi$. By Theorem 1.6, the semigroup $(T_t \otimes Id_E)_{t \geqslant 0}$ is R-analytic on $L^p(M, E)$. By Proposition 5.2, these two results imply that A_p actually admits a bounded $H^\infty(\Sigma_\theta)$ functional calculus for some $0 < \theta < \frac{\pi}{2}$, as expected.

Finally, it is not hard to check that the above arguments work as well with $(Id_{B(\ell^2)} \otimes T_t)_{t \geqslant 0}$ in the place of $(T_t)_{t \geqslant 0}$. thus A_p actually has a completely bounded functional calculus for some $0 < \theta < \frac{\pi}{2}$. ■

Now, we give some natural examples to which the results of this paper can be applied.

Noncommutative Poisson semigroup. Let $n \geqslant 1$ be an integer. Here $\mathbb{F}_n$ denotes a free group with n generators denoted by $g_1, \ldots, g_n$. Any $g \in \mathbb{F}_n$ has a unique decomposition of the form

$$g = g_{i_1}^{k_1} g_{i_2}^{k_2} \cdots g_{i_l}^{k_l},$$

where $l \geqslant 0$ is an integer, each i_j belongs to $\{1, \ldots, n\}$, each k_j is a non zero integer, and $i_j \neq i_{j+1}$ if $1 \leqslant j \leqslant l-1$. The case when $l = 0$ corresponds to the unit element $g = e_{\mathbb{F}_n}$. By definition, the length of g is defined as

$$|g| = |k_1| + \cdots + |k_l|.$$

This is the number of factors in the above decomposition of g. For any nonnegative real number $t \geqslant 0$, we have a normal unital completely positive selfadjoint map

$$\begin{array}{rccc} T_t : & \mathrm{VN}(\mathbb{F}_n) & \longrightarrow & \mathrm{VN}(\mathbb{F}_n) \\ & \lambda(g) & \longmapsto & e^{-t|g|}\lambda(g). \end{array}$$

These maps define a w*-semigroup $(T_t)_{t \geqslant 0}$ called the noncommutative Poisson semigroup (see [JMX] for more information). In [Arh, remark following Proposition 5.5], it is implicitly said that $(T_t)_{t \geqslant 0}$ is $QWEP$-dilatable. Similarly, we can show each T_t is $QWEP$-factorizable.

q-Ornstein-Uhlenbeck semigroup. We use the notations of [BKS]. Suppose $-1 \leqslant q < 1$. Let H be a real Hilbert space and let $(a_t)_{t\geqslant 0}$ be a strongly continuous semigroup of contractions on H. For any $t \geqslant 0$, let $T_t = \Gamma_q(a_t)$. Then $(T_t)_{\geqslant 0}$ is a w*-semigroup of normal unital completely positive maps preserving the trace on the von Neumann algebra $\Gamma_q(H)$.

In the case where $a_t = e^{-t} I_H$, the semigroup $(T_t)_{\geqslant 0}$ is the so-called q-Ornstein-Uhlenbeck semigroup.

Using [Ric], [Nou] and the result [KW, Theorem 10.11] of dilation of strongly continuous semigroups of contractions on a Hilbert space it is not hard to see that we obtain examples of $QWEP$-dilatable semigroups of $QWEP$-factorizable maps.

We pass to Schur multipliers.

Theorem 5.3 *Let $(T_t)_{t\geqslant 0}$ be a w^*-semigroup self-adjoint contractive Schur multipliers on $B\big(\ell^2_I\big)$. Suppose $1 < p, q < \infty$ and $0 < \theta < 1$. Let E be an operator space such that $E = \big(OH(I), F\big)_\theta$ for some index set I and for some $OUMD_q$-operator space F such that $\frac{1}{p} = \frac{1-\theta}{2} + \frac{\theta}{q}$. We let $-A_p$ be the generator of the strongly continuous semigroup $(T_t \otimes Id_E)_{t\geqslant 0}$ on $S^p_I(E)$. Then for some $0 < \theta < \frac{\pi}{2}$, the operator A_p has a bounded $H^\infty(\Sigma_\theta)$ functional calculus.*

Proof : Arguing as in the proof of [Arh, Corollary 4.3], we may reduce the general case to the unital and completely positive case. The proof for semigroups of unital completely positive Schur multipliers, using [Arh, Proposition 5.5], is similar to the one of Theorem 1.4. ■

Remark 5.4 *The results of this paper lead to properties of some square functions, see [LM, section 7] for more information.*

Acknowledgment. The author would like to express his gratitude to Christian Le Merdy for some useful advices. He also thank Hun Hee Lee and Quanhua Xu for some valuable discussions on Proposition 2.2 and Éric Ricard for an interesting conversation.

References

[AnD] C. Anantharaman-Delaroche. On ergodic theorems for free group actions on noncommutative spaces. Probab. Theory Related Fields 135 (2006), no. 4, 520–546.

[Arh] C. Arhancet. On Matsaev's conjecture for contractions on noncommutative L^p-spaces. To appear in Journal of Operator theory. Preprint, arXiv:1009.1292.

[BeG] E. Berkson and T. Gillespie. Spectral decompositions and harmonic analysis on UMD Banach spaces. Studia Math. 112 (1994), 13–49.

[Beu] A. Beurling. On analytic extension of semigroups of operators. J. Functional Analysis 6 (1970), 387–400.

[Bou] J. Bourgain Vector-valued singular integrals and the H^1-BMO duality. Probability theory and harmonic analysis (Cleveland, Ohio, 1983), 1–19, Monogr. Textbooks Pure Appl. Math., 98, Dekker, New York (1986).

[BKS] M. Bożejko, B. Kümmerer and R. Speicher. q-Gaussian processes: non-commutative and classical aspects. Comm. Math. Phys. 185 (1997), no. 1, 129–154.

[ClP] P. Clement, B. de Pagter, F. A. Sukochev and H. Witvliet. Schauder decomposition and multiplier theorems. Studia Math. 138 (2000), no. 2, 135–163.

[Con] A. Connes. Classification of injective factors. Cases II_1, II_∞, III_λ, $\lambda \neq 1$. Ann. of Math. (2) 104 (1976), no. 1, 73–115.

[CwR] M. Cwikel and S. Reisner. Interpolation of uniformly convex Banach spaces. Proc. Amer. Math. Soc. 84 (1982), no. 4, 555–559.

[DJT] J. Diestel, H. Jarchow and A. Tonge. Absolutely summing operators. Cambridge Studies in Advanced Mathematics, 43. Cambridge University Press (1995).

[DH] M. Dostál and D. Hadwin. An alternative to free entropy for free group factors. International Workshop on Operator Algebra and Operator Theory (Linfen, 2001). Acta Math. Sin. (Engl. Ser.) 19 (2003), no. 3, 419–472.

[ER] E. Effros and Z.-J. Ruan. Operator spaces. Oxford University Press (2000).

[Fac] S. Fackler. Regularity of Semigroups via the Asymptotic Behaviour at Zero. Preprint, arXiv:1203.5498.

[HaM] U. Haagerup and M. Musat. Factorization and dilation problems for completely positive maps on von Neumann algebras. Comm. Math. Phys. 303 (2011), no. 2, 555–594.

[HJX] U. Haagerup, M. Junge and Q. Xu. A reduction method for noncommutative L_p-spaces and applications. Trans. Amer. Math. Soc. 362 (2010), no. 4, 2125–2165.

[Haa] M. Haase. The functional calculus for sectorial operators. Operator Theory: Advances and Applications, 169. Birkhäuser Verlag (2006).

[Har] A. Harcharras, Asma. On some "stability" properties of the full C^*-algebra associated to the free group F_∞. Proc. Edinburgh Math. Soc. (2) 41 (1998), no. 1, 93–116.

[Jun1] M. Junge. Fubini's theorem for ultraproducts of noncommmutative L_p-spaces. Canad. J. Math. 56 (2004), no. 5, 983–1021.

[Jun2] M. Junge. Fubini's theorem for ultraproducts of noncommmutative L_p-spaces II. Preprint.

[HJX] U. Haagerup, M. Junge and Q. Xu. A reduction method for noncommutative L_p-spaces and applications. Trans. Amer. Math. Soc. 362 (2010), no. 4, 2125–2165.

[JMX] M. Junge, C. Le Merdy and Q. Xu. H^∞ functional calculus and square functions on noncommutative L^p-spaces. Astérisque No. 305 (2006).

[JuP] M. Junge and J. Parcet. The norm of sums of independent noncommutative random variables in $L_p(l_1)$. J. Funct. Anal. 221 (2005), no. 2, 366–406.

[KaW] N. J. Kalton, L. Weis. The H^∞-calculus and sums of closed operators. Math. Ann. 321 (2001), no. 2, 319–345.

[KW] P. C. Kunstmann and L. Weis. Maximal L_p-regularity for parabolic equations, Fourier multiplier theorems and H^∞-functional calculus. pp. 65-311 in Functional analytic methods for evolution equations, Lect. Notes in Math. 1855, Springer (2004).

[LM] C. Le Merdy. Square functions, bounded analytic semigroups, and applications. Perspectives in operator theory, 191–220, Banach Center Publ., 75, Polish Acad. Sci., Warsaw (2007).

[Mus] M. Musat. On the operator space UMD property for noncommutative L_p-spaces. Indiana Univ. Math. J. 55 (2006), no. 6, 1857–1891.

[Nou] A. Nou. Asymptotic matricial models and $QWEP$ property for q–Araki–Woods algebras. J. Funct. Anal. 232 (2006), no. 2, 295–327.

[Oza] N. Ozawa. About the $QWEP$ conjecture. Internat. J. Math. 15 (2004), no. 5, 501–530.

[Pau] V. Paulsen. Completely bounded maps and operator algebras. Cambridge Univ. Press (2002).

[Pis1] G. Pisier. Sur les espaces de Banach K-convexes (French). Seminar on Functional Analysis, 1979–1980 (French), Exp. No. 11, 15 pp., École Polytech., Palaiseau (1980).

[Pis2] G. Pisier. Holomorphic semigroups and the geometry of Banach spaces. Ann. of Math. (2) 115 (1982), no. 2, 375–392.

[Pis3] G. Pisier. Regular operators between non-commutative L_p-spaces. Bull. Sci. Math. 119 (1995), no. 2, 95–118.

[Pis4] G. Pisier. The operator Hilbert space OH, complex interpolation and tensor norms. Mem. Amer. Math. Soc. 122 (1996), no. 585.

[Pis5] G. Pisier. Non-commutative vector valued L_p-spaces and completely p-summing maps. Astérisque, 247 (1998).

[Pis6] G. Pisier. Introduction to operator space theory. Cambridge University Press, Cambridge (2003).

[Qiu] Y. Qiu. On the OUMD property for the column Hilbert space C. Preprint, arXiv:1107.4941.

[Ric] É. Ricard. A Markov dilation for self-adjoint Schur multipliers. Proc. Amer. Math. Soc. 136 (2008), no. 12, 4365–4372.

[Ste] E. M. Stein. Topics in harmonic analysis related to the Littlewood-Paley theory. Annals of Mathematics Studies, No. 63 Princeton University Press, Princeton, N.J.; University of Tokyo Press, Tokyo (1970).

Laboratoire de Mathématiques, Université de Franche-Comté, 25030 Besançon Cedex, France
cedric.arhancet@univ-fcomte.fr